

Biococc

Toltrazuril

Oral solution

For veterinary use

Target species

Broilers, breeding poultry, turkeys, geese, ducks

Pharmacotherapeutic group

Antiprotozoal agent

ATC vet code: **QP51BC01** – Toltrazuril

Composition

1 ml contains: **Toltrazuril 25 mg.**

Excipients: **Polyethylene glycol up to 1 ml.**

Description

Colorless, transparent liquid with a characteristic odor.

Presentation

- 100 ml in a polyethylene bottle
 - 1 L in a polyethylene bottle
 - 1 ml in a glass ampoule
-

Pharmacological properties

Active against coccidia (*Eimeria acervulina*, *Eimeria brunetti*, *Eimeria necatrix*, *Eimeria mitis*, *Eimeria adenoides*, *Eimeria meleagridis*, *Eimeria anseris*, *Eimeria truncata*), including strains resistant to other anticoccidial products.

Biococc is moderately toxic for warm-blooded animals. When administered at recommended doses and at doses increased 10-fold, it does not cause side effects or

complications in poultry and does not inhibit the formation of immunity against coccidiosis.

Indications

Used for prevention and treatment of coccidiosis in broilers, breeding poultry, turkeys, geese and ducks.

Method of administration and dosage

Biococc is administered with drinking water for **2 days** at a dose of **7 mg toltrazuril per 1 kg body weight** of poultry (**28 ml of 2.5% Biococc solution per 100 kg body weight**), which is equivalent to:

- **1 ml of 2.5% product dissolved in 1 liter of drinking water**, given to poultry to drink for **24 hours**,
or
- **3 ml of product per 1 liter of drinking water** for **8 hours**, for **2 consecutive days**.

In severe cases of the disease, the treatment course is repeated after **5 days**.

The medicinal solution of Biococc remains active for **48 hours**.

Biococc is used in combination with feed additives and vitamins. Shake well before use.

Side effects

When used at recommended doses according to indications, no side effects were observed.

Contraindications

Not to be used in laying hens.

Special instructions

The product is compatible with feed additives, vitamins and other poultry medicinal products.

When working with Biococc, follow personal hygiene rules and safety precautions for handling veterinary medicines.

In case of contact with skin or mucous membranes, rinse with a stream of water.

Withdrawal period

Poultry may be slaughtered for meat **8 days** after the last administration of the product.

If forced slaughter occurs earlier than this period, the meat may be used for production of meat-and-bone meal.

Not to be used in laying hens!

Storage conditions

Store in a dry place protected from light. The product must be stored tightly closed in the original manufacturer packaging at a temperature not exceeding **25°C**.

Keep out of reach of children.

Shelf life: **2 years**.

Manufacturer

LLC **Biotecs**, Tbilisi, Yumashvili St. **#8**