

Calfostonic

Oral powder

Composition

1 kg contains:

Vitamin A – 600,000 IU

Vitamin D3 – 200,000 IU

Vitamin B1 – 100 mg

Vitamin B2 – 200 mg

Vitamin B6 – 10 mg

Vitamin B12 – 1.1 mg

Vitamin E – 75 mg

Vitamin K3 – 25 mg

Niacinamide – 1.25 g

Calcium pantothenate – 500 mg

Choline chloride – 25 g

DL-Methionine – 10 g

L-Lysine – 2.5 g

Gentian root – 5 g

L-Carnitine – 3 g

Sodium glutamate – 7.5 g

Sodium chloride – 24.6 g

Magnesium carbonate – 50.0 g

Manganese (sulfate) – 480 mg

Zinc (sulfate) – 504 mg

Iron (sulfate) – 600 mg

Copper (sulfate) – 125 mg

Cobalt (carbonate) – 100 mg

Potassium iodide – 124 mg

Calcium carbonate – 443.9 g

Dicalcium phosphate – 300 g

Sodium selenite – 80 mg

Excipients q.s. up to 1 kg

Characteristics

Calfostonic is a complex mixture of mineral salts, water- and fat-soluble vitamins, trace elements, amino acids, non-specific growth factors, appetite stimulants, tonics, and aromatic additives.

When administered to animals, it corrects acute and chronic deficiency conditions and provides essential nutrients necessary for proper growth and full development. As a result, milk, meat, and wool production can reach levels consistent with each animal's genetic potential.

Indications

Calfostonic is used for treatment and prevention of:

- **Acute and chronic deficiencies**
 - **Anorexia**
 - **Unbalanced nutrition**
 - **Recovery periods**
 - **Rickets**
 - **Poor development in animals**
 - **Situations requiring tonics and appetite stimulants**
-

Contraindications

Not described.

Adverse Reactions

Not described.

Target Species

Cattle, sheep, goats, poultry, pigs, and horses.

Dosage and Administration

Group treatment

All species:

20 kg per ton of feed for 10–20 days.

Poultry:

20–30 g per kg body weight, mixed in feed, for 10–20 days.

Individual treatment

Cattle:

Adults: 100–200 g per animal once daily for 15–20 days.

Calves: 40–50 g per animal once daily for 15–20 days.

Pigs and horses:

During pregnancy: 60–100 g per animal once daily for 15–20 days.

Young animals: 10–15 g per animal once daily for 15–20 days.

Withdrawal Period

Not required.

Storage Conditions

Store in a cool place protected from light at 5°C–25°C.

Shelf Life

4 years.

Special Warnings

Overdose is unlikely due to the composition.

Presentation

1 kg polyethylene bag.

Keep out of reach of children.

For veterinary use only.

Available without veterinary prescription.

Manufacturer

Biotecs, Georgia