

## **Ichthyol Ointment 20%**

**(Ichthyol Ointment 20%)**

**Ointment for external and topical use in veterinary medicine**

---

### **Composition**

**1 g of ointment contains:**

- **Ichthyol – 200 mg**
- 

### **Description**

**Black or dark brown ointment with a characteristic odor.**

---

### **Pharmacological Properties**

**The active substance, ichthyol, has a complex composition containing aromatic compounds and ammonium salts of sulfonic acids.**

**It exhibits:**

- **Antimicrobial action**
- **Anti-inflammatory effect**
- **Antiparasitic activity**
- **Keratoplastic effect**
- **Local anesthetic effect**

**When applied to damaged skin areas, ichthyol ointment causes mild irritation of sensitive nerve endings, followed by a decrease in their sensitivity and improvement of tissue trophism.**

**The ointment has a positive effect on the skin and subcutaneous tissues. In the area of inflammation, it regulates vascular and capillary tone, restores blood circulation, reduces exudation, promotes resorption of infiltrates, and accelerates tissue regeneration.**

---

## **Indications for Use**

**Used in animals for skin conditions such as:**

- **Open wounds**
  - **Purulent wounds**
  - **Burns**
  - **Eczema**
  - **Dermatitis**
  - **Scars and cracks**
- 

## **Adverse Reactions**

**No adverse reactions have been observed.**

**Hypersensitivity to the components of the preparation may occur.**

---

## **Contraindications**

**None established.**

---

## **Target Species**

**All animal species.**

---

## **Dosage and Administration**

**The ointment is applied locally by rubbing into the affected area and surrounding tissues, or applied in a thin layer to clean white cloth or gauze.**

**Dressings and applications are performed 2–3 times daily until recovery.**

---

## **Overdose**

**No overdose symptoms have been identified.**

---

### **Specificity of Action**

**No specific effects have been identified upon first application or withdrawal.**

---

### **Pregnancy and Lactation**

**The product may be used during pregnancy and lactation.**

---

### **Withdrawal Period**

**0 days.**

---

### **Precautions**

**Avoid contact of the ointment with eyes, nose, mouth, and teats of lactating animals.**

---

### **Storage Conditions**

**Store in a dry place protected from light at a temperature not exceeding 20°C.  
Keep out of reach of children.**

---

### **Shelf Life**

**3 years.**

---

### **Presentation**

- **20 g in polyethylene container**
  - **0.5 kg in polyethylene container**
- 

### **Pharmacotherapeutic Group**

**Antiseptic agent.**

**ATC vet code: QD08AX – Other antiseptics and disinfectants.**

---

**Manufacturer**

**LLC “Biotecs”**

**Georgia, Tbilisi**

**8 Iumashev Street**