

## **Iodoform Powder**

**Powder for external use**

**For all animal species**

---

### **Composition**

**1 g of powder contains the following active substances:**

- **Iodoform – 0.011 g**
  - **Boric acid – 0.149 g**
  - **Oxytetracycline hydrochloride – 0.04 g**
  - **Streptocide – 0.05 g**
- 

### **Description**

**Powder ranging in color from light yellow to yellow, with a characteristic specific odor.**

---

### **Pharmacological Properties**

**The antiseptic action of the product is based on the properties of its active ingredients.**

**Iodoform – Upon contact with the wound surface, active iodine is released and gradually decomposes, ensuring a prolonged antibacterial effect.**

**Boric acid – An antiseptic substance with insecticidal properties. Promotes protein coagulation, including enzymatic proteins. Disrupts cell membrane permeability, suppresses phagocytosis, and inhibits bacterial growth and development.**

**Oxytetracycline hydrochloride – A broad-spectrum antibiotic from the tetracycline group. Exhibits bacteriostatic activity and is active against many Gram-positive and Gram-negative bacteria, rickettsiae, spirochetes, large viruses, and some protozoa.**

**Streptocide – A short-acting sulfonamide with bacteriostatic properties. Effective against streptococci, meningococci, gonococci, Escherichia coli, and other microorganisms. Due to antagonism with para-aminobenzoic acid, it blocks folic acid synthesis in microbial cells, thereby inhibiting microbial growth and development.**

---

## **Indications for Use**

**Used for the treatment of:**

- **Infected wounds**
- **Post-operative wounds**
- **Ulcers**
- **Skin eruptions**
- **Moist eczema**
- **Dermatitis**
- **Tendovaginitis**
- **Phlegmon**
- **Lymphadenitis**
- **Slow-healing wounds**

---

## **Adverse Reactions**

**When used according to instructions, no adverse reactions or complications (allergic skin reactions, local irritation) have been observed.**

---

## **Contraindications**

**None established.**

---

## **Target Species**

**Animals (all species).**

---

## **Dosage and Administration**

**The product is applied locally by sprinkling onto the affected area.**

**Before application, clean the affected area from dirt and remove necrotic tissue.**

**Apply a thin layer 1–2 times daily.**

**Prevent animals from licking the treated area.**

**The need for repeated treatment depends on clinical (surgical) findings and severity of the lesion.**

---

#### **Interaction with Other Medicinal Products**

**No data available regarding interactions with other products.**

---

#### **Overdose**

**No data available.**

---

#### **Pregnancy and Lactation**

**External use of iodoform powder during pregnancy and lactation is not restricted.**

---

#### **Withdrawal Period**

**Meat and milk: 0 days.**

---

#### **Special Instructions**

**For target species**

**If atypical symptoms occur after starting treatment, consult a veterinarian.**

**For persons administering the product**

**When handling the product, follow personal hygiene and veterinary safety regulations.**

**Eating, drinking, and smoking during handling are prohibited.**

**After work, wash hands thoroughly with warm water and soap.**

**Persons hypersensitive to product components should avoid direct contact.**

**If accidental contact with skin or mucous membranes occurs, rinse with running water.**

**In case of allergic reaction or accidental ingestion, seek medical attention immediately and present the product label or instructions.**

---

#### **Storage Conditions**

**Store in a dry place protected from light at a temperature not exceeding 25°C.**

---

#### **Shelf Life**

**2 years.**

---

#### **Presentation**

**50 g powder for external use in a polyethylene container.**

---

#### **Pharmacotherapeutic Group**

**Antibiotic, dermatological medicinal product.**

**ATCvet code: QD06AA53 – Oxytetracycline, combinations.**

---

**For veterinary use only.**

**Dispensed by veterinary prescription.**

**Keep out of reach of children.**

---

#### **Manufacturer**

**LLC “Biotecs”**

**Georgia, Tbilisi**

**8 Iumashev Street**