

Ivomectin-F

Ivermectin + Clorsulon

Injectable Solution For Veterinary Use Only

Description

Ivomectin-F is a broad-spectrum antiparasitic injectable solution containing a combination of ivermectin and clorsulon. The product is effective against trematodes, nematodes, and various ectoparasites. It is used for the treatment and control of parasitic diseases in cattle, sheep, and goats.

Composition

Each 1 ml injectable solution contains:

- Ivermectin – 10 mg
- Clorsulon – 100 mg

Target Species

Cattle, sheep, and goats.

Indications for Use

The product is indicated for the treatment of mixed infestations caused by trematodes, nematodes, and external parasites.

Trematodes

- Liver flukes (*Fasciola hepatica* and *Fasciola gigantica*) – immature stages.

Gastrointestinal Nematodes

- *Ostertagia* spp.
- *Haemonchus placei*
- *Trichostrongylus* spp.
- *Cooperia* spp.
- *Oesophagostomum radiatum*
- *Nematodirus* spp.
- *Strongyloides papillosus*
- *Chabertia* spp.
- *Bunostomum phlebotomum*

Lung Nematodes

- Dictyocaulus viviparus (adult stages and fourth-stage larvae)

External Parasites

- Hypoderma bovis
- Hypoderma lineatum
- Ticks
- Biting lice
- Mange mites
- Sucking lice

Dosage and Administration

For subcutaneous injection only.

Cattle, Sheep and Goats

- 1 ml per 50 kg body weight.

Contraindications

- Do not administer intravenously or intramuscularly.
- Do not use in animals with known hypersensitivity to the active ingredients.

Adverse Reactions

- Temporary discomfort may occur following subcutaneous administration.
- Mild swelling or pain may occur at the injection site.
- Hypersensitivity reactions may rarely occur.

Special Warnings

- Administer only by subcutaneous injection.
- To avoid local reactions, doses exceeding 10 ml should be divided between several injection sites.
- For optimal control of Hypoderma lineatum, treatment is recommended immediately after the end of the fly season.
- Use with caution during periods of larval migration.

Interactions with Other Medicinal Products

Do not use concurrently with:

- Ketoconazole

- Itraconazole
- Cyclosporine
- Calcium channel blockers
- Diethylcarbamazine

Pregnancy and Lactation

The product is safe for use in pregnant and lactating animals.

Overdose

Significant overdose of ivermectin may result in:

- Depression
- Apathy
- Ataxia
- Recumbency
- Death in severe cases

Treatment is symptomatic.

Withdrawal Period

- Meat: 49 days.

Milk: Do not use in dairy animals producing milk for human consumption or in animals within two months before calving.

Presentation

100 ml injectable solution supplied in a clear glass vial sealed with a rubber stopper, aluminum cap, and tamper-evident plastic flip-off cap. Packed in a cardboard box.

Storage Conditions

- Store in a dry place protected from light.
- Store below 30°C.
- Keep out of reach of children.

Shelf life: 3 years.

Shelf life after first opening: 28 days.

Manufacturer

Biotecs LLC

8 Iumashevi Street, Tbilisi, Georgia.