

Instructions for Use – Fir Oil Ointment

Dosage Form: Ointment for external use, used in veterinary medicine.

Pharmacotherapeutic Group: Antiseptic, anti-inflammatory agent.

Composition (per 1 g of ointment):

Active substance: Fir oil – 70 mg

Excipients: Vaseline – 930 mg

Description: Light yellow ointment with a characteristic odor.

Packaging: 20 g ointment in polyethylene container.

Pharmacological Properties:

When applied to the skin, the ointment diffuses through sebaceous and sweat glands, from where it is continuously absorbed. It causes vasodilation, improves blood supply, accelerates resorption during inflammation, promotes growth of granulation tissue and wound epithelialization.

Indications:

Used for treatment of wounds, burns, eczema, dermatitis, bursitis, and traumatic tissue injuries.

Administration and Dosage:

Apply a thin layer of the ointment onto clean gauze and bandage the wound. Dressings are changed twice daily until recovery. The ointment may also be applied directly to the affected area with gentle rubbing. In case of injuries and mastitis, warm compresses with ointment are recommended.

Adverse Reactions:

No adverse reactions or complications have been observed in animals.

Contraindications:

Not established.

Storage Conditions:

Store protected from light at a temperature not exceeding 25°C.

Shelf life: 2 years.

Manufacturer:

LLC "Biotecsi", Georgia, Tbilisi, Iumashev #8.