

Sultrim – Oral tablets for veterinary use

Pharmacotherapeutic Group: Systemic antimicrobial, antibacterial agent.

Target Species: Cattle, small ruminants, carnivorous animals, swine, poultry, rabbits.

Composition:

Each tablet contains:

- Sulfadimidine sodium – 400 mg
 - Trimethoprim – 80 mg
-

Indications for Use

Sultrim tablets are used for the treatment of infectious diseases in domestic animals caused by microorganisms sensitive to sulfamethoxazole and trimethoprim, including:

- Upper and lower respiratory tract infections;
- Gastrointestinal tract diseases;
- Genitourinary infections;
- Secondary infections during viral diseases;
- Skin and soft tissue infections.

Cattle, small ruminants, carnivores:

Diarrhea, colisepticemia, streptococcal polyarthrititis, salmonellosis, pasteurellosis, pneumonia, staphylococcal abscesses.

Swine:

Colibacillosis, atrophic rhinitis, salmonellosis, pasteurellosis, toxoplasmosis, haemophilus pleuropneumonia, erysipelas.

Poultry:

Coccidiosis, pasteurellosis, colisepticemia, salmonellosis, staphylococcosis, infectious coryza.

Rabbits:

Coccidiosis, salmonellosis, staphylococcosis.

Dosage and Administration

Cattle, small ruminants, carnivores, swine, rabbits:

Administer mixed with feed, water, or milk at a dose of:

- 1 tablet (480 mg) per 15 kg body weight, every 12 hours for 3–5 days.

Poultry:

- Chicks up to 1 week of age: 1/4 tablet (120 mg) dissolved in 2 L of drinking water.
- Broiler and layer chicks (1–8 weeks): 1/2 tablet (240 mg) dissolved in 2 L of water or mixed with 1 kg of feed.
- Laying hens (over 20 weeks): 1/2 tablet (240 mg) dissolved in 1 L of drinking water or 3/4 tablet (360 mg) mixed with 1 kg of feed.
- Turkeys (5–20 weeks): 1/2 tablet (240 mg) per 1 L of water.

When administering with water, dissolve the required amount of the product first in a small quantity of water (100–150 ml) until a uniform suspension is obtained, then add to the remaining water until fully dissolved.

Treatment in all animal species should continue for 2 additional days after clinical signs disappear.

Pharmaceutical Form

90 tablets in a polyethylene bottle.

Storage Conditions

Store in a dry place protected from light at a temperature not exceeding 20°C.

Shelf Life

3 years.

Manufacturer

Biotecs, Georgia.