

**Product Name: Phistoclozanid (PHISTOCLOZANID)**

**Active Substance: Oxyclozanide**

**Group: Antiparasitic, anthelmintic**

**Dosage Form: Tablet, bolus**

---

### **Description**

Phistoclozanid is used for the prevention and treatment of liver, gastric, and intestinal helminth infections in large and small ruminants (cows, calves, sheep, goats).

Its active substance, oxyclozanide, acts on the nervous system of parasites, ensuring their complete elimination.

---

### **Indications**

Fasciolosis, paramphistomosis, moniezirosis, and other helminthic diseases.

---

### **Dosage and Administration**

#### **Cattle and calves:**

- Moniezirosis and paramphistomosis: 1 tablet (750 mg) per 50 kg body weight.
- Fasciolosis: 1 tablet (750 mg) per 75 kg body weight.

#### **Sheep and goats:**

- Same dosage, administered with feed.
  - After 2 hours, provide water to induce mild laxative effect to eliminate dead parasites.
- 

### **Withdrawal Period**

- Meat: 14 days
- Milk: 3 days

Do not use during the first and last trimester of pregnancy.

- ◆ For consultation regarding use, visit: [www.roqi.ge](http://www.roqi.ge)
- 

### Packaging

- Tablets: 500 mg, 750 mg (90 and 300 tablets)
  - Boluses: 1500 mg (30 boluses), 3000 mg (60 boluses)
- 

### Storage Conditions

Store in a dry place protected from light at a temperature below 25°C.

Store in a dry place protected from light at 20°C.

---

### Shelf Life

3 years.

---

### Manufacturer

Biotecs, Georgia  
8 Iumashevi Street